

Using linear electrodynamic quadrupole trap for drop pattern formation and nanoparticle aggregates production from drying colloidal microdroplets of suspension

Justice Archer^{a,*}, Mariusz Woźniak^a, Tomasz Wojciechowski^{a,b}, Maciej Kolwas^a, Daniel Jakubczyk^a, Gennadiy Derkachov^a, and Krystina Kolwas^a

^a*Institute of Physics, Polish Academy of Sciences, Aleja Lotnikow 32/46, PL-02668 Warsaw, Poland*

^b*International Research Centre MagTop, Aleja Lotnikow 32/46, PL-02668 Warsaw, Poland*

*Presenting author (archer@ifpan.edu.pl)

Electrodynamic quadrupole trapping of microparticles provides a contactless approach for optical measurements of microdroplets of different compositions, aerosols, and dust grains [1,2]. The use of the linear electrodynamic quadrupole trap (LEQT) [3] can provide unsupported solvent drying, well-defined aggregate morphology evolutions from colloidal droplets of suspension as well the possibility to deposit (“soft-landing”) the final dry microproducts on a substrate and analyze them offline by scanning electron microscopy (SEM) [4,5].

In this presentation, we will demonstrate advancements in the use of the LEQT to produce various nanoparticle aggregates of interesting structural properties from drying microdroplets of colloidal suspension. The final products were carefully deposited on a substrate and analyzed offline with scanning electron microscopy. Highly-ordered aggregates of silica nanoparticles, aggregates of gold and silica as well as aggregates of silica with crystallized sodium dodecylsulfate (SDS) will be presented. Additionally, we will present SEM analysis of intermediate drying drop deposits and the manner in which SDS alter these deposits/patterns with time and SDS concentration. Our ability to deposit intermediate drop patterns and complex dry aggregates of different morphologies with unique properties using the LEQT and offline SEM analysis is desirable in many applications.

References

- [1] Hart M. B, *et al.* Optical measurements from single levitated particles using a linear electrodynamic quadrupole trap. *Appl. Opt.* 2015;54:174–81.
- [2] Ray A. K, *et al.* Resonance-based light scattering techniques for investigation of microdroplet processes. *Faraday Discuss.* 2007;137:85–98.
- [3] Paul W. Electrodynamic traps for charged and neutral particles. *Rev. Mod. Phys.* 1990;62.
- [4] Woźniak M, *et al.* Formation of highly ordered spherical aggregates from drying microdroplets of colloidal suspension. *Langmuir* 2015;31.
- [5] Archer J, *et al.* Evolution of radius and light scattering properties of single drying microdroplets of colloidal suspension. *J. Quant. Spectrosc. Radiat. Transfer* 2017;202.

Preferred mode of presentation: Oral/Poster